

3. Ideal Solution

〈Solution Equilibrium of Gas〉

In a solution equilibrium state of a gas into water, $A(g) \rightleftharpoons A(aq)$:

$$\mu(g) = \mu(aq) \quad (3.1)$$

For an ideal gas and an ideal solution [from (0.6) and (0.7)]:

$$-\Delta_{\text{sol}}G^\circ = \mu^\circ(g) - \mu^\circ(aq) = RT \ln K_{\text{sol}} \quad (3.2)$$

$$\text{where } K_{\text{sol}} \equiv \frac{b/b^\circ}{p/p^\circ}$$

The Henry's law constant, K_H , is often defined in the reverse way as,

$$p = K_H b \quad (3.3)$$

$$K_H = K_{\text{sol}}^{-1} \quad (3.4)$$

Exercise 3.1

1) Compute the solution equilibrium constant of CO_2 into water, $K_{\text{sol}} = (b/b^\circ)/(p/p^\circ)$, from the following standard Gibbs energies.

	$\Delta_f G^\circ(298 \text{ K}) / \text{kJ mol}^{-1}$
$\text{CO}_2(g)$	-394.4
$\text{CO}_2(aq)$	-386.0

2) Compare the result with the well established Henry's law constant, $K_H = 30.1 \text{ bar kg mol}^{-1}$.

Solution to exercise 3.1

1) $\Delta_{\text{sol}}G^\circ = -386.0 - (-394.4) = 8.4 \text{ kJ mol}^{-1}$.

$$K_{\text{sol}} = \exp(-\Delta_{\text{sol}}G^\circ / RT) = \exp[-8400 / (8.3145 \times 298)] = 3.37 \times 10^{-2} \text{ (mol kg}^{-1} \text{ bar}^{-1}\text{)}$$

2) K_H from thermo data, $K_H = K_{\text{sol}}^{-1} = 29.7 \text{ bar kg mol}^{-1}$, agrees well with the given $K_H = 30.1$.

〈Heat of Solution and Temperature Dependence〉

The eq. (3.2) can be rewritten as,

$$K_{\text{sol}} = \exp\left(\frac{\Delta_{\text{sol}}S^\circ}{R}\right) \exp\left(-\frac{\Delta_{\text{sol}}H^\circ}{RT}\right) \quad (3.5)$$

For $\Delta H < 0$ (exothermic) $K \uparrow$ as $T \downarrow$, while for $\Delta H > 0$ (endothermic) $K \uparrow$ as $T \uparrow$.

(Le Chatelier's principle)

Exercise 3.2

The enthalpy of solution of H_2 into water is $\Delta_{\text{sol}}H^\circ = -4.2 \text{ kJ mol}^{-1}$ and the solution equilibrium constant is $K_{\text{sol}} = 7.81 \times 10^{-4} \text{ mol kg}^{-1} \text{ bar}^{-1}$ at 298 K. Estimate K_{sol} at 10 °C (283 K).

Solution to exercise 3.2

$$\frac{K_{\text{sol}, 283}}{K_{\text{sol}, 298}} = \exp\left[-\frac{\Delta_{\text{sol}}H^\circ}{R} \left(\frac{1}{283} - \frac{1}{298}\right)\right] = \exp\left[-\frac{-4.2 \times 1000}{8.3145} \left(\frac{1}{283} - \frac{1}{298}\right)\right] = 1.094$$

Then, $K_{\text{sol}, 283} = 7.81 \cdot 10^{-4} \times 1.094 = 8.54 \times 10^{-4} \text{ mol kg}^{-1} \text{ bar}^{-1}$.

cf.) experimental = $8.72 \times 10^{-4} \text{ mol kg}^{-1} \text{ bar}^{-1}$

K_{sol} increases \uparrow as T decreases \downarrow . (consistent with $\Delta H < 0$; exothermic)

(alternative solution)

$$\Delta_{\text{sol}}G^\circ = -RT \ln K_{\text{sol}} = -8.3145 \cdot 298 \ln(7.81 \times 10^{-4}) = 17.73 \text{ kJ mol}^{-1}.$$

$$\Delta_{\text{sol}}S^\circ = (\Delta_{\text{sol}}H^\circ - \Delta_{\text{sol}}G^\circ) / T = (-4.2 - 17.73) \times 1000 / 298 = -73.59 \text{ J K}^{-1} \text{ mol}^{-1}.$$

By assuming constant $\Delta_{\text{sol}}H^\circ$ and $\Delta_{\text{sol}}S^\circ$,

$$\Delta_{\text{sol}}G^\circ(283 \text{ K}) = -4.2 - 283 \times (-73.59) / 1000 = 16.63 \text{ kJ mol}^{-1}$$

$$K_{\text{sol}} = \exp(-\Delta_{\text{sol}}G^\circ / RT) = \exp[-16630 / (8.3145 \times 283)] = 8.5_2 \times 10^{-4} \text{ mol kg}^{-1} \text{ bar}^{-1}.$$

Exercise 3.3Estimate the solubility S (mol kg⁻¹) of AgCl(s) at 50 °C from followings.

(298 K)	$\Delta_f H^\circ / \text{kJ mol}^{-1}$	$S_m^\circ / \text{J K}^{-1} \text{ mol}^{-1}$
AgCl(s)	-127.1	96.2
Ag ⁺ (aq)	105.6	72.7
Cl ⁻ (aq)	-167.2	56.5

Solution to exercise 3.3

$$\Delta_{\text{sol}}H^\circ = (105.6 - 167.2) - (-127.1) = 65.5 \text{ kJ mol}^{-1},$$

$$\Delta_{\text{sol}}S^\circ = (72.7 + 56.6) - 96.2 = 33.0 \text{ J K}^{-1} \text{ mol}^{-1}.$$

$$\text{at } 50 \text{ }^\circ\text{C}, S = \exp[(-65500 / 8.3145 / 323 + 33.0 / 8.3145) / 2] = 3.68 \times 10^{-5} \text{ mol kg}^{-1}$$

$$\text{at } 25 \text{ }^\circ\text{C}, S = \underline{1.31 \times 10^{-5}} \text{ (} \rightarrow \text{ exercise 1.3)}$$

cf.) experimental: 3.81×10^{-5} (50 °C), 1.35×10^{-5} (25°C)Solubility S increases \uparrow as T increases \uparrow . (consistent with $\Delta H > 0$; endothermic)**(Activity)**Except for the very dilute solution such as in exercise 3.3, the solution equilibrium of the electrolyte, $\text{AB(s)} \rightleftharpoons \text{A}^+(\text{aq}) + \text{B}^-(\text{aq})$, should be written with the mean activity coefficient, γ , as,

$$\begin{aligned} -\Delta_r G^\circ &= \mu^\circ[\text{AB(s)}] - \mu^\circ[\text{A}^+] - \mu^\circ[\text{B}^-] \\ &= RT \ln \left(\frac{\gamma b[\text{A}^+]}{b^\circ} \frac{\gamma b[\text{B}^-]}{b^\circ} \right) \end{aligned} \quad (3.6)$$

Exercise 3.41) Assuming the ideal solution, calculate the solubility S (mol kg⁻¹) of KCl(s) from the followings.

	$\Delta_f G^\circ(298 \text{ K}) / \text{kJ mol}^{-1}$
KCl(s)	-409.1
K ⁺ (aq)	-283.3
Cl ⁻ (aq)	-131.2

2) Evaluate the mean activity coefficient, γ , for the saturated KCl solution by using the measured solubility $S = 4.769 \text{ mol kg}^{-1}$.**Solution to exercise 3.4**

$$1) \Delta_r G^\circ = (-283.3) + (-131.2) - (-409.1) = -5.4 \text{ kJ mol}^{-1}$$

$$b[\text{A}^+(\text{aq})]b[\text{B}^-(\text{aq})] = \exp(-\Delta_r G^\circ / RT) = \exp[5.4 \times 1000 / (8.3145 \times 298)] = 8.84 \text{ mol}^2 \text{ kg}^{-2}$$

$$S = (8.84)^{1/2} = \underline{2.97} \text{ mol kg}^{-1}$$

$$2) \gamma = 2.97 / 4.769 = \underline{0.623}$$

Trend of solubility S against T for small $|\Delta H|$ dissolution process often disturbed (looks reversed) by T -dep of γ .