

# Homogeneous Kinetics

## 5. Elementary Reactions

### ⟨Rate Equation⟩

Rate of reaction:  $m\text{A} + n\text{B} + \dots \rightarrow i\text{X} + j\text{Y} + \dots$

$$v = -\frac{1}{m} \frac{d[\text{A}]}{dt} = -\frac{1}{n} \frac{d[\text{B}]}{dt} = \dots = \frac{1}{i} \frac{d[\text{X}]}{dt} = \frac{1}{j} \frac{d[\text{Y}]}{dt} = \dots \quad (5.1)$$

[A], [B], ...: concentrations of A, B, ...

Rate equation

$$v = k[\text{A}]^m[\text{B}]^n \dots \quad (5.2)$$

$k$ : rate constant

### Exercise 5.1

- 1) Write the rate equation for an irreversible reaction,  $\text{A} \rightarrow \text{B}$  (rate const. =  $k_1$ ), with respect to A, and solve the differential equation (rate equation) for the initial condition,  $[\text{A}] = [\text{A}]_0$  at  $t = 0$ .
- 2) Write the rate equation for an irreversible reaction,  $2\text{A} \rightarrow \text{B}$  (rate const. =  $k_2$ ), with respect to A, and solve it for the initial condition,  $[\text{A}] = [\text{A}]_0$  at  $t = 0$ .

#### Solution to exercise 5.1

1) rate equation:  $(v =) -\frac{d[\text{A}]}{dt} = k_1[\text{A}]$ . solution:  $[\text{A}] = [\text{A}]_0 \exp(-k_1 t)$ .

2) rate equation:  $(v =) -\frac{1}{2} \frac{d[\text{A}]}{dt} = k_2[\text{A}]^2$ . solution:  $\frac{1}{[\text{A}]} = \frac{1}{[\text{A}]_0} + 2k_2 t$  or  $[\text{A}] = \frac{[\text{A}]_0}{1 + 2k_2 t [\text{A}]_0}$ .

### ⟨Elementary Reaction⟩

≡ Minimum step of reaction that obeys eq. (5.2)

Examples:

1)  $\text{H}_2 + \text{Br}_2 \rightarrow 2\text{HBr}$ :  $v = \frac{1}{2} \frac{d[\text{HBr}]}{dt} \propto \frac{[\text{H}_2][\text{Br}_2]^{3/2}}{[\text{Br}_2] + c[\text{HBr}]}$  NO

· complex sequence of reactions:  $\text{Br}_2 \rightarrow 2\text{Br}$ ,  $\text{Br} + \text{H}_2 \rightleftharpoons \text{HBr} + \text{H}$ ,  
 $\text{H} + \text{Br}_2 \rightarrow \text{HBr} + \text{Br}$ , etc.

2)  $\text{OH} + \text{H}_2 \rightarrow \text{H}_2\text{O} + \text{H}$ :  $v = \frac{d[\text{H}_2\text{O}]}{dt} = k[\text{OH}][\text{H}_2]$  YES

### Exercise 5.2

Argue whether the reaction,  $\text{H}_2 + \text{I}_2 \rightarrow 2\text{HI}$ , is an elementary reaction or not, from the following measurements for the initial rate of formation at 600 K.

exp. #	$[\text{H}_2]$ / mol m <sup>-3</sup>	$[\text{I}_2]$ / mol m <sup>-3</sup>	$d[\text{HI}]/dt _{t=0}$ / mol m <sup>-3</sup> s <sup>-1</sup>
#1	0.72	0.51	0.175
#2	0.72	1.02	0.350
#3	1.44	1.02	0.700

#### Solution to exercise 5.2

from #1 & #2,  $d[\text{HI}]/dt \propto [\text{I}_2]$ ; from #2 & #3,  $d[\text{HI}]/dt \propto [\text{H}_2]$ . So,  $v \propto [\text{H}_2][\text{I}_2]$  and this reaction CAN be an elementary reaction.

\* Eq. (5.2) may be accidentally satisfied. (but this is really an elementary reaction.)

### Consecutive Reactions

Rate equations for the consecutive reactions,  $A \xrightarrow{k_1} B \xrightarrow{k_2} C$ , with respect to [A], [B] and [C]

$$\frac{d[A]}{dt} = -k_1[A], \quad \frac{d[B]}{dt} = k_1[A] - k_2[B], \quad \frac{d[C]}{dt} = k_2[B] \quad (5.3)$$

Solutions for  $k_1 \neq k_2$  and for the initial conditions,  $[A] = [A]_0$  and  $[B] = [C] = 0$  at  $t = 0$


$$[A] = [A]_0 \exp(-k_1 t), \quad [B] = \frac{k_1 [A]_0}{k_1 - k_2} \{ \exp(-k_2 t) - \exp(-k_1 t) \},$$

$$[C] = [A]_0 - [A] - [B] \quad (5.4)$$

#### Exercise 5.3

- 1) Complete the following table of solution (5.4) for  $k_1 = 5$ ,  $k_2 = 1$  and  $[A]_0 = 1$  and plot concentrations [A], [B] and [C].


$t$	[A]	[B]	[C]
0	1	0	0
0.1	0.61	0.37	0.02
0.2	0.37	0.56	0.07
0.4	0.14	0.67	0.19
0.7	0.03	0.58	0.39
1.1	0	0.41	0.59
1.6	0	0.25	0.75
2.3	0	0.13	0.87
3	0	0.06	0.94



- 2) Describe which parts of the time-profile of [B] represent  $k_1$  and  $k_2$ .

- 3) Complete the following table of solution (5.4) for  $k_1 = 1$ ,  $k_2 = 5$  and  $[A]_0 = 1$  and plot concentrations.

$t$	[A]	[B]	[C]
0	1	0	0
0.1	0.90	0.07	0.03
0.2	0.82	0.11	0.07
0.4	0.67	0.13	0.20
0.7	0.50	0.12	0.38
1.1	0.33	0.08	0.59
1.6	0.20	0.05	0.75
2.3	0.10	0.03	0.87
3	0.05	0.01	0.94



- 4) Describe which parts of the time-profile of [B] represent  $k_1$  and  $k_2$ .

#### Solution to exercise 5.3

- As shown in the figure to the right.
- [B] rises with  $k_1$  ( $\tau_1 = k_1^{-1} = 0.2$ ) and decays with  $k_2$  ( $\tau_2 = k_2^{-1} = 1$ ).
- As shown in the figure to the right.
- [B] rises with  $k_2$  ( $\tau_2 = k_2^{-1} = 0.2$ ) and decays with  $k_1$  ( $\tau_1 = k_1^{-1} = 1$ ).

\* Exactly the same solution for [C]!

\* Similar (same except for the height) solution for [B]!

\* For [B],  $k_1$  &  $k_2$  look reversed when  $k_2 > k_1$ .

